Medications Quiz (3rd Year)

What is AF?		
	nticoagulant used in AF to minimise the risk of clo	ots?
What is the drug used t	to reverse the effects of the above medication?	
NovorapidLevemirHumulin	lin short, medium or long acting?	
	tions should not be given before theatre?	
Omeprazole Bisoprolol	Clopidogrel Zomorph	Levothyroxine Domperidone
What is the reason for	this?	
	between Phosphate Sandoz and Sando K ?	
What observation can f	Fludrocortisone help to treat and where is it store	ed?

What should I do!?

pressure is 82/40. You look through their previous observations to find that this is sometimes normal for them. The patient is prescribed 80mg Furosemide . Do you give the medication? If yes or no, please give an explanation as to your decision.
One of your patients is prescribed Meropenem and they have a Penicillin allergy. What should you do?
You are caring for a patient who has come back from theatre at 16:00 following a DHS. You are administering medications at 18:00 and your patient is due 4,500 units of Tinzaparin . Before administering the mediation, what two things should you check?
You are working on nights and you are performing the controlled drug checks. When checking Oxycodone you find that there should be 28 tablets. However, you can only find 14, therefore there is a strip missing. What actions should you take?
You are caring for a patient with type 1 diabetes and they are prescribed 10 units of Insulatard in the morning. You take their blood sugar before breakfast and it is 3.8. The patient states they do not feel like any breakfast this morning and may have something to eat later. Do you administer or omit the insulin?